



For more information about the Organization of Teratology Information Specialists or to find a service in your area, call (866) 626-6847 or visit us online at: www.OTISpregnancy.org.

Benzodiazepines and Pregnancy

This sheet talks about the risks that exposure to benzodiazepines can have during pregnancy. With each pregnancy, all women have a 3% to 5% chance of having a baby with a birth defect. This information should not take the place of medical care and advice from your health care provider.

What is a benzodiazepine?

Benzodiazepines are medications used to treat anxiety, sleeplessness, seizures, muscle spasms, and alcohol withdrawal. Valium[®] (diazepam), Xanax[®] (alprazolam), Klonopin[®] (clonazepam), and Ativan[®] (lorazepam) are a few examples of benzodiazepines, but there are many others. While it is best to study medicines individually, benzodiazepines are often studied together during pregnancy.

Should I stop taking my benzodiazepine once I find out I'm pregnant?

No. You should always talk to your doctor before making any changes in your medication. If you suddenly stop taking your medication you may have withdrawal and we don't know what effect withdrawal might have on a pregnancy. Your doctor can help you decide if the benefit of taking the medicine outweighs any possible risk to your pregnancy.

I've heard that benzodiazepines can cause birth defects like cleft lip and palate. Is this true?

Some early studies in animals and humans suggested a slight increase in the risk for cleft lip and/or cleft palate if a benzodiazepine was taken during the first trimester. Since these early reports, there have been studies and reviews that have not

supported an association between benzodiazepines and cleft lip or palate or birth defects in general. It is generally felt that exposure to a benzodiazepine does not significantly increase the risk for birth defects.

Can taking benzodiazepines cause other pregnancy problems?

Two studies have suggested a higher rate of preterm deliveries and low birth weight in infants when women take benzodiazepines during pregnancy. However, a third study did not find these risks, so more research is needed to accurately answer this question. It is possible that other factors and not the medicine were responsible for these findings.

I need to continue taking my benzodiazepine medication. Will it cause any harmful effects in my baby after birth?

If you are taking a benzodiazepine near the time of delivery, your baby may have withdrawal symptoms such as not being able to control body temperature, difficulty breathing, muscle weakness, irritability, crying, sleep disturbances, tremors, and jitteriness. It is important that you inform your obstetrician and your baby's pediatrician so extra care can be provided should your baby need it. These symptoms resolve over a period of time as the drug leaves the baby's system and are not expected to have any long-term effects.

Will taking a benzodiazepine have any effect on my baby's behavior and development?

Some studies in animals have suggested an effect on behavior in exposed offspring. However, since animals do not always predict the effects in humans, no conclusions can be made. Presently, there are no well-done, long-term studies looking at children exposed to benzodiazepines during pregnancy. However, these drugs have been on the market for more than 40 years and there has been no evidence to suggest that they have long-term harmful effects on the child's brain or development.

Someone told me not to take my benzodiazepine with my allergy medication diphenhydramine. Why not?

A single human report and animal data have suggested that the combination of temazepam and diphenhydramine may increase risk for stillbirth or death shortly after birth. It is presently unknown if this interaction will occur with all benzodiazepines. To be safe, it is best not to take diphenhydramine while you are taking a benzodiazepine.

Can I take benzodiazepines while I am breastfeeding?

Some benzodiazepines are not recommended during breastfeeding because they stay in the body a long time and can potentially cause sedation in a breastfed infant. If a benzodiazepine is needed during breastfeeding, it is best to use one that is removed from the body rapidly (such as lorazepam). When using a benzodiazepine during breastfeeding, watch your baby for sleepiness, low energy, or poor suckling which may be signs your baby is getting too much of the drug. If any of these symptoms are seen, discuss them with your pediatrician promptly.

What if the father of the baby takes a benzodiazepine?

Since men do not share a blood connection with the baby, an increase risk of birth defects or pregnancy complications is not expected when the father of the baby takes a benzodiazepine. For more information about a father's exposures and pregnancy, please see the OTIS fact sheet [*Paternal Exposures and Pregnancy*](#).

July 2010.
Copyright by OTIS.
Reproduced by permission.



Selected References:

- Barry WS, et al. SM. 1987. Exposure to benzodiazepines in utero. *Lancet* 1:1436-37.
- Calderon-Margalit R, et al. 2009. Risk of preterm delivery and other adverse perinatal outcomes in relation to maternal use of psychotropic medications during pregnancy. *Am J Obstet Gynecol*. 201(6):579.e1-8.
- Czeizel AE. 1988. Lack of evidence of teratogenicity of benzodiazepine drugs in Hungary. *Reprod Toxicol* 1:183-88.
- Czeizel AE, Toth M 1998. Birth weight, gestational age and medications during pregnancy. *Int J Obstet Gynaecol Obstet* 60:245-49.
- Eros E, et al. 2002. A population-based case-control teratologic study of nitrazepam, medazepam, tofisopam, alprazolam, and clonazepam treatment during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 101:147-54.
- Kargas GA et al 1985 Perinatal mortality due to interaction of diphenhydramine and temazepam. *N Eng J Med* 313:1417-18
- Laegreid L, et al.1992. The effect of benzodiazepines on the fetus and the newborn. *Neuropediatrics* 23:18-23.
- Laegreid L, et al.1992. Neurodevelopment in late infancy after prenatal exposure to benzodiazepines—a prospective study. *Neuropediatrics* 23:60-67.
- McElhatton PR, 1994. The effects of benzodiazepine use during pregnancy and lactation. *Reprod Toxicol* 8(6):461-75.
- Ornoy A, et al.1998. Is benzodiazepine use during pregnancy really teratogenic? *Reprod Toxicol* 12:511-5.
- Sanchis A, et al. 1991. Adverse effects of maternal lorazepam on neonates. *DICP Ann Pharmacother* 25(10):1137-38.
- Shiono PH, et al. 1984. Oral clefts and diazepam use during pregnancy. *N Engl J Med* 311:919-20.
- Weber LWD. 1985. Benzodiazepines in pregnancy—academic debate or teratogenic risk? *Biol Res Pregnancy* 6(4):151-67.
- Wikner BN, et al. 2007. Use of benzodiazepines and benzodiazepine receptor agonists during pregnancy: maternal characteristics. *Pharmacoepidemiol Drug Saf* 16(9):988-994.
- Wikner BN, et al. 2007. Use of benzodiazepines and benzodiazepine receptor agonists during pregnancy: neonatal outcome and congenital malformations *Pharmacoepidemiol Drug Saf* 16(11):1203-1210.

*If you have questions about the information on this fact sheet or other exposures during pregnancy, call **OTIS** at **1-866-626-6847**.*